

Hindcast & Nowcast of Currents in the Santa Barbara

Lie-Yauw Oey

Program in Atmospheric & Oceanic Sciences

Sayre Hall, Forrestal Campus

Princeton University

Princeton, NJ 08544

Phone: (609) 258-5971 Fax: (609) 258-2850

Email: lyo@splash.princeton.edu

Award #:N00014-95-1-0809

LONG-TERM GOALS

To develop a high-resolution, verifiable nowcast and forecast model for the Santa Barbara Channel (SBC) by combining basic understanding of the circulation processes through theory and observations, numerical techniques and data assimilation.

OBJECTIVES

To test the model using inputs from realistic wind and density fields for Dec/97 through May/98, and use the results to study the momentum balance in the channel.

APPROACH

Oey's (1996) nested-grid model for the Southern California Bight (SCB) and SBC (Figure 1) was extended to hindcast and nowcast currents using up-to-date windstresses and CalCofi temperature and salinity (T and S) data during the recent El Nino winter and spring, from Dec/97 through May/98. Windstresses were calculated by combining hourly NDBC wind in the vicinity of the channel with historical, monthly COADS wind over the outer region away from the channel. Historical T and S fields were used to initialize the model, which were then supplemented with a simple data-assimilation scheme that nudged the modeled T/S to CalCofi data of Dec/97 through Apr/98 (Courtesy of Tom Hayward, Scripps Inst. Oceanogr.) The assimilation of CalCofi data introduced waters with temperatures that are approximately 2° warmer than climatology in the Southern California Bight.

WORK COMPLETED

Model code development was completed.

RESULTS

Figure 2 shows examples of surface temperature (left panel) and free-surface elevation (right panel), with surface (left) and 100-m (right) velocity vectors superimposed, for (a) Jan/15, (b) Feb/14, (c) Mar/16 and (d) Apr/15, 1998. Advection of the El Nino warm water by strong poleward coastal current can be seen in January. A momentum balance analysis (Oey et al., 1998) shows that at this time, poleward pressure gradient caused by sea-level tilt is balanced primarily by the tendency term $\partial v / \partial t + \partial v^2 / \partial y$. From February through May, the equatorward windstress intensified, and one begins to see the onset of equatorward coastal currents in Figure 2b. The near-shore waters also became cooler

Report Documentation Page			<i>Form Approved OMB No. 0704-0188</i>					
<p>Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p>								
1. REPORT DATE 1998	2. REPORT TYPE	3. DATES COVERED 00-00-1998 to 00-00-1998						
4. TITLE AND SUBTITLE Hindcast & Nowcast of Currents in the Santa Barbara			5a. CONTRACT NUMBER					
			5b. GRANT NUMBER					
			5c. PROGRAM ELEMENT NUMBER					
6. AUTHOR(S)			5d. PROJECT NUMBER					
			5e. TASK NUMBER					
			5f. WORK UNIT NUMBER					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Princeton University, Program in Atmospheric & Oceanic Sciences, Princeton, NJ, 08544			8. PERFORMING ORGANIZATION REPORT NUMBER					
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)					
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)					
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited								
13. SUPPLEMENTARY NOTES See also ADM002252.								
14. ABSTRACT								
15. SUBJECT TERMS								
16. SECURITY CLASSIFICATION OF: <table border="1" style="display: inline-table; vertical-align: middle;"> <tr> <td style="padding: 2px;">a. REPORT unclassified</td> <td style="padding: 2px;">b. ABSTRACT unclassified</td> <td style="padding: 2px;">c. THIS PAGE unclassified</td> </tr> </table>			a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified						

brought about by equatorward advection and upwelling. The upwelling process can be seen clearly in March (Figure 2c), where cool waters are found off the central California coast, and warmer waters are 'pushed back' in the SCB. At this time, a cyclone also begins to form in the western part of the channel, with waters cooler than 11.5° and sea-level of about -4cm at its center. By April, a significant reversal of the nearshore currents, from being equatorward to poleward, can be seen to originate in the SCB (Figure 2d). Warm waters, and together with it higher sea-level also, begin to propagate poleward along the coast. This phenomenon can be attributed to forcing by the equatorward weakening of the windstress curl that occurs south of Pt.Conception in the SCB (Oey,1996,'98).

IMPACT/APPLICATIONS

Sensitivity experiments were also conducted (details in Oey et al., 1998). Of these, the most important finding is that the SBC circulation depends on the relative intensity of windstresses west and east of the channel. This is because the poleward pressure gradient in the channel tends to be balanced by equatorward windstress. Since friction is small, any imbalance is taken up by the Coriolis acceleration of the cross-channel flow. The acceleration is equatorward if the wind is uniformly intense west and east, and poleward if the wind is much weaker in the east. The former condition induces southward cross-channel flow and corresponds to the observed 'Flood East' or 'Upwelling' scenario as defined by Harms and Winant (1998), while the latter northward cross-channel flow and to the 'Cyclonic' or 'Relaxation' scenario. These results are of practical value as they identify key forcing parameters that will guide in further refinements of the model's skill in the channel.

TRANSITIONS

Results in the form of IEEE binary files have been submitted to the Navy (contact: SAIC) for analysis in conjunction with Apr/May 1998 field experiments in the channel.




Figure 1 Santa Barbara Channel and Southern California Bight locator map and model domain with topography in meters. For computational efficiency, maximum model depth is set to 2000 m. Dashed rectangle area is the sub-region shown in Figure 2.

RELATED PROJECTS

The research is in part supported by a grant from The Mineral Management Service (Contract # DOI-10094286; Program Manager: David Brown) via Scripps Institute of Oceanography, so that I work closely with Clinton Winant (SIO) on observational data aspect, and also with Dong-Ping Wang (SUNY), on modeling and data-assimilation aspects.

REFERENCES & PUBLICATIONS

Harms, S. and Winant, C.D., 1998: Characteristics patterns of the circulation in the Santa Barbara Channel. *J. Geophys. Res.* 103: 3041-3065.

Oey, L.-Y., 1996: Flow around a coastal bend: a model of the Santa Barbara Channel eddy. *J. Geophys. Res.*, 101, 16,667-16,682.

Oey, L.-Y., 1998: A Forcing Mechanism for the Poleward Flow off the Southern California Coast. Submitted to *J. Geophys. Res.*

Oey, L.-Y., C. Winant, M. Hendershott and T. Hayward, 1998: Momentum Balance from a Hindcast and Nowcast Model of Currents in the Santa Barbara Channel. Manuscript. (Also, available as a report to ONR, October, 1998).

Figure 2 The one-day averaged surface temperature (left panel) and free-surface elevation contours (right panel), with surface (left) and 100-m (right) velocity vectors superimposed, for (a) Jan/15, (b) Feb/14, (c) Mar/16 and (d) Apr/15, 1998. Five-day averaged wind stresses are also plotted as vectors with thick arrows at four locations. For clarity, only a portion of the model domain (enclosed in dashed rectangle in Figure 1) focusing on the Santa Barbara Channel and vicinity is shown.




Figure 2a




Figure 2b




Figure 2c




Figure 2d